

## PRIOR AUTHORIZATION POLICY

**POLICY:** Enzyme Replacement Therapy – Kanuma Prior Authorization Policy

- Kanuma™ (sebelipase alfa intravenous infusion – Alexion)

**REVIEW DATE:** 04/06/2022

---

### OVERVIEW

Kanuma, a human lysosomal acid lipase (LAL), indicated for the treatment of patients with a diagnosis of **LAL deficiency**.<sup>1</sup> It is produced in the egg white of genetically engineered chickens via recombinant DNA technology. LAL catalyzes the breakdown of cholesteryl esters to free cholesterol and fatty acids, and the breakdown of triglycerides to glycerol and free fatty acids.

### Disease Overview

LAL deficiency is a rare lysosomal storage disorder characterized by absent or deficient LAL activity leading to the accumulation of cholesterol and triglycerides in the liver and other organs.<sup>2,3</sup> Patients with LAL deficiency often have dyslipidemias, cardiovascular disease and progressive liver disease.<sup>2</sup> The disorder has a heterogeneous presentation ranging from a rapidly progressive form occurring in infants which leads to death in the first year of life, to a childhood/adult-onset form with milder signs and symptoms. Almost all patients with childhood/adult-onset LAL deficiency have hepatomegaly with elevated liver transaminases and have an increased risk of developing fibrosis and cirrhosis.<sup>3</sup> The diagnosis of LAL deficiency is established by demonstrating deficient LAL activity in leukocytes, fibroblasts, or liver tissue, or by genetic testing.<sup>2,3</sup>

### POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Kanuma. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Kanuma as well as the monitoring required for adverse events and long-term efficacy, approval requires Kanuma to be prescribed by or in consultation with a physician who specializes in the condition being treated.

**Automation:** None.

### RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Kanuma is recommended in those who meet the following criteria:

#### FDA-Approved Indication

- 1. Lysosomal Acid Lipase Deficiency.** Approve for 1 year if the patient meets the following criteria (A and B):
  - A)** The diagnosis is established by one of the following (i or ii):
    - i.** Patient has a laboratory test demonstrating deficient lysosomal acid lipase activity in leukocytes, fibroblasts, or liver tissue; OR
    - ii.** Patient has a molecular genetic test demonstrating lysosomal acid lipase gene mutation; AND
  - B)** Kanuma is prescribed by or in consultation with a geneticist, endocrinologist, a metabolic disorder sub-specialist, or a physician who specializes in the treatment of lysosomal storage disorders.

04/06/2022

© 2022. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

**CONDITIONS NOT RECOMMENDED FOR APPROVAL**

Coverage of Kanuma is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

**REFERENCES**

1. Kanuma™ intravenous infusion [prescribing information]. Cheshire, CT: Alexion; November 2021.
2. Reiner Z, Guardamagna O, Nair D, et al. Lysosomal acid lipase deficiency – an under-recognized cause of dyslipidaemia and liver dysfunction. *Atherosclerosis*. 2014;235:21-30.
3. Erwin AL. The role of sebelipase alfa in the treatment of lysosomal acid lipase deficiency. *Ther Adv Gastroenterol*. 2017;10:553-562.