

PRIOR AUTHORIZATION POLICY

- POLICY:** Gonadotropin-Releasing Hormone Antagonists – Oriahnn Prior Authorization Policy
- Oriahnn™ (elagolix, estradiol, and norethindrone acetate capsules; elagolix capsules – AbbVie)

REVIEW DATE: 04/26/2023

OVERVIEW

Oriahnn, an oral gonadotropin-releasing hormone (GnRH) receptor antagonist with added estrogen and progestin therapy, is indicated for the **management of heavy menstrual bleeding associated with uterine leiomyomas (fibroids) in premenopausal women.**¹ Limitation of Use: Use should be limited to 24 months due to the risk of continued bone loss which may not be reversible.¹

Disease Overview

Uterine fibroids (leiomyomas) are benign tumors. They are the most frequent gynecologic benign disease.² Fibroids can be asymptomatic or cause symptoms; symptoms generally present as abnormal (heavy) uterine bleeding or pelvic pain/pressure. Heavy menstrual bleeding can cause associated problems, such as iron deficiency anemia. The actual prevalence of uterine fibroids is difficult to ascertain since many patients are asymptomatic, but it is estimated that fibroids can be detected in up to 80% of women by 50 years of age.³

Guidelines

Oriahnn is addressed in the American College of Obstetrician and Gynecologists guidelines on the management of symptomatic uterine leiomyomas (2021).⁴ Medical treatment options for uterine leiomyomas include agents that address only bleeding symptoms, such as GnRH antagonists, levonorgestrel-releasing intrauterine devices, contraceptive steroids, and tranexamic acid. Agents that reduce both bleeding and leiomyoma size include GnRH agonists and selective progesterone receptor modulators (SPRMs). SPRMs are not approved in the U.S. for the treatment of uterine leiomyomas. An oral GnRH antagonist, such as Oriahnn or Myfembree, can be considered for the treatment of abnormal uterine bleeding related to leiomyomas for up to 2 years. The hormonal add-back therapy is indicated to offset the hypoestrogenic effects of the product.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Oriahnn. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days. Because of the specialized skills required for evaluation and diagnosis of patients treated with Oriahnn as well as the monitoring required for adverse events and long-term efficacy, approval requires Oriahnn to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

04/26/2023

© 2022. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Oriahnn is recommended in those who meet the following criteria:

FDA-Approved Indication

1. **Uterine Fibroids (Leiomyomas).** Approve for up to 24 months if the patient meets the following criteria (A, B, C, D, E, F, and G):

Note: Approve for **up to** 24 months. For example, a patient who has already received 6 months of treatment with Oriahnn should be approved for a duration of 18 months.

- A) Patient is ≥ 18 years of age; AND
- B) Patient is PREmenopausal (before menopause); AND
- C) Patient is experiencing heavy menstrual bleeding associated with the uterine fibroids; AND
- D) Uterine fibroids have been confirmed by a pelvic ultrasound, including transvaginal ultrasonography or sonohysterography; hysteroscopy; or magnetic resonance imaging; AND
- E) Patient has tried at least one other therapy for the medical management of heavy menstrual bleeding; AND

Note: Examples of therapy for the medical management of heavy menstrual bleeding includes: combination estrogen-progestin contraceptives (oral tablets, vaginal ring, transdermal patch), levonorgestrel-releasing intrauterine systems [e.g. Mirena, Liletta], oral progesterone (e.g., medroxyprogesterone acetate), depo-medroxyprogesterone injection, tranexamic acid tablets.

- F) Patient has **not** previously received a continuous regimen of 24 months or longer of therapy with Oriahnn or Myfembree; AND
- G) The medication is prescribed by or in consultation with an obstetrician-gynecologist or a health care practitioner who specializes in the treatment of women's health.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Oriahnn is not recommended in the following situations:

1. **Heavy Menstrual Bleeding not associated with Uterine Fibroids.**

Oriahnn has been shown to be effective in reducing heavy menstrual bleeding only in women with uterine fibroids.¹

2. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Oriahnn™ co-packaged capsules [prescribing information]. North Chicago, IL: AbbVie; August 2021.
2. Neri M, Melis G, Giancane E, et al. Clinical utility of elagolix as an oral treatment for women with uterine fibroids: A short report on the emerging efficacy data. *Int J Womens Health*. 2019;11:535-546.
3. De La Cruz MS, Buchanan EM. Uterine Fibroids: Diagnosis and Treatment. *Am Fam Physician*. 2017;95(2):100-107.
4. American College of Obstetricians and Gynecologists. ACOG Practice Bulletin. Management of Symptomatic Uterine Leiomyomas. June 2021. Available at: <https://www.acog.org/clinical/clinical-guidance/practice-bulletin/articles/2021/06/management-of-symptomatic-uterine-leiomyomas>. Accessed on April 20, 2023.

