

## PRIOR AUTHORIZATION POLICY

**POLICY:** Hyperlipidemia – Nexletol Prior Authorization Policy

- Nexletol® (bempedoic acid tablets – Esperion)

**REVIEW DATE:** 04/26/2023; selected revision 05/03/2023 and 01/17/2024

---

### OVERVIEW

Nexletol, an adenosine triphosphate-citrate lyase inhibitor, is indicated as an adjunct to diet and statin therapy for the treatment of primary hyperlipidemia in adults with the following:<sup>1</sup>

- **Atherosclerotic cardiovascular disease (ASCVD)** for those who require additional lowering of low-density lipoprotein cholesterol (LDL-C).
- **Heterozygous familial hypercholesterolemia (HeFH)** for those who require additional lowering of LDL-C.

The safety and effectiveness have not been established in pediatric patients.<sup>1</sup>

### Clinical Efficacy

CLEAR Outcomes was a randomized, double-blind, placebo-controlled trial involving 13,970 adults, 18 to 85 years of age who were unable or unwilling to take statins due to unacceptable adverse events among those who had, or were at high risk for, CV disease.<sup>2</sup> Those eligible had to report being statin intolerant. Patients were assigned to receive Nexletol or placebo. Use of statins at very low doses were permitted, as well as other lipid lowering therapies (e.g., ezetimibe, bile acid sequestrants, fibrates). The mean patient age was 65 years. In total, 70% of patients had a previous CV event (secondary prevention population) whereas 30% of patients were categorized as being in the primary prevention group. At baseline, 22.7% of patients were utilizing a statin and 11.5% were on ezetimibe. The mean LDL-C at baseline was 139 mg/dL. The median follow-up was 40.6 months. The mean LDL-C level after 6 months of treatment with Nexletol was 107 mg/dL for the Nexletol group vs. 136 mg/dL for placebo. The primary endpoint (death from CV causes, nonfatal myocardial infarction [MI], nonfatal stroke, or coronary revascularization) occurred in 11.7% of patients in the Nexletol group vs. 13.3% in the placebo group (P = 0.004). The composite of death from CV causes, nonfatal stroke, or nonfatal MI occurred in 8.2% of patients given Nexletol vs. 9.5% of patients in the placebo group (P = 0.006).

### Guidelines

Many guidelines are available regarding the treatment of patients with dyslipidemia which include the management of HeFH and ASCVD.<sup>3-10</sup> For patients with elevated LDL-C, statins are the cornerstone of therapy and recommended first-line to be used at maximally tolerated doses due to the established benefits regarding the reduction of CV risks. Atorvastatin 40 mg to 80 mg once daily (QD) and rosuvastatin 20 mg to 40 mg QD are considered high-intensity statins as they achieve LDL-C lowering of  $\geq 50\%$ .

- The **American College of Cardiology Expert Consensus Decision Pathway on the Role of Non-Statins Therapies** for LDL-Cholesterol Lowering in the Management of ASCVD Risk (2022) make several recommendations; Nexletol is addressed.<sup>3</sup> For adults with clinical ASCVD at very high risk (e.g., patients with major ASCVD events, HeFH, diabetes) who are on statin therapy for secondary prevention, the general goal is  $\geq 50\%$  LDL-C reduction and an LDL-C < 55 mg/dL (of non-HDL-C < 85 mg/dL) with maximally tolerated statin therapy. If the above goals are not achieved, the initial non-statin agents recommended include ezetimibe and/or a proprotein convertase subtilisin kexin type 9 (PCSK9) monoclonal antibody (i.e., Repatha or Praluent). Nexletol can be considered after these therapies.

04/26/2023

© 2023. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

- The **American Association of Clinical Endocrinologists and American College of Endocrinology** has guidelines regarding the management of dyslipidemia and the prevention of CV disease (2020).<sup>7</sup> Nexletol is mentioned is cited as an option for intensification of therapy after use of standard agents such as high-intensity/moderate-intensity statins.
- The **International Lipid Expert Panel** published a position paper in 2023 on use of Nexletol in the management of lipid disorders and CV risk.<sup>10</sup> One recommendation is that in patients with statin intolerance, Nexletol monotherapy, or in combination with ezetimibe and other non-statin drugs is recommended to enable patients to achieve therapeutic goals. In primary prevention, Nexletol may be considered for patients at high and very high CV risk who despite optimally maximally tolerated doses of statins and ezetimibe, are not achieving target LDL-C levels.
- The **American Heart Association (AHA)/American College of Cardiology** guidelines on the management of blood cholesterol (2018) define ASCVD as acute coronary syndrome, those with a history of MI, stable or unstable angina or coronary or other revascularizations, stroke, transient ischemic attack, or peripheral arterial disease.<sup>4,5</sup> An LDL-C < 70 mg/dL is recommended for most patients with ASCVD to reduce CV risk. Guidelines and reviews have recognize that patients with an elevated coronary artery calcium or calcification score (e.g.,  $\geq 300$  Agatston units) are at an increased risk of CV events.<sup>8,11-14</sup>
- **The AHA** published a scientific statement regarding familial hypercholesterolemia (2015).<sup>9</sup> Key points are that the condition may start early (in childhood or adolescence) and is noted by LDL-C levels  $\geq 190$  mg/dL. Premature CV disease can result. Diagnosis can be confirmed by genetic testing. The Dutch Lipid Clinic Network criteria and Simon Broome criteria may also be used which incorporate cholesterol levels, family history, clinical findings, and physical manifestations. Aggressive lipid-lowering therapy is recommended to achieve LDL-C reductions of at least 50%.

## POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Nexletol. All approvals are provided for the duration noted below. A patient who has previously met Initial Therapy criteria for Nexletol for the requested indication under the Coverage Review Department and is currently receiving Nexletol is only required to meet continuation of therapy criteria (i.e., currently receiving therapy). If past criteria has not been met under the Coverage Review Department and the patient is currently receiving Nexletol, or is restarting Nexletol, Initial Therapy criteria must be met.

**Automation:** None.

## RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Nexletol is recommended in those who meet the following criteria:

### FDA-Approved Indications

1. **Atherosclerotic Cardiovascular Disease.\*** Approve for 1 year if the patient meets ONE of the following (A or B):
  - A) **Initial Therapy.** Approve if the patient meets all of the following (i, ii, and iii):
    - i. Patient is  $\geq 18$  years of age; AND
    - ii. Patient has had one of the following conditions or diagnoses (a, b, c, d, e, or f):
      - a) A previous myocardial infarction or a history of an acute coronary syndrome; OR
      - b) Angina (stable or unstable); OR
      - c) A past of stroke or transient ischemic attack; OR
      - d) Coronary artery disease; OR

04/26/2023

© 2023. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

- e) Peripheral arterial disease; OR
  - f) Patient has undergone a coronary or other arterial revascularization procedure in the past; AND  
Note: Examples include coronary artery bypass graft surgery, percutaneous coronary intervention, angioplasty, and coronary stent procedures.
- iii. Patient meets one of the following (a or b):
- a) Patient meets all of the following [(1), (2), and (3)]:
    - (1) Patient has tried one high-intensity statin therapy (i.e., atorvastatin  $\geq$  40 mg daily; rosuvastatin  $\geq$  20 mg daily [as a single entity or as a combination product]); AND
    - (2) Patient has tried one high-intensity statin above along with ezetimibe (as a single-entity or as a combination product) for  $\geq$  8 continuous weeks; AND
    - (3) Low-density lipoprotein cholesterol level after this treatment regimen remains  $\geq$  70 mg/dL; OR
  - b) Patient has been determined to be statin intolerant by meeting one of the following [(1) or (2)]:
    - (1) Patient experienced statin-related rhabdomyolysis; OR  
Note: Rhabdomyolysis is statin-induced muscle breakdown that is associated with markedly elevated creatine kinase levels (at least 10 times the upper limit of normal), along with evidence of end organ damage which can include signs of acute renal injury (noted by substantial increases in serum creatinine [Scr] levels [ $a \geq 0.5$  mg/dL increase in Scr or doubling of the Scr] and/or myoglobinuria [myoglobin present in urine]); OR
    - (2) Patient meets all of the following [(a), (b), and (c)]:
      - (a) Patient experienced skeletal-related muscle symptoms; AND  
Note: Examples of skeletal-related muscle symptoms include myopathy (muscle weakness) or myalgia (muscle aches, soreness, stiffness, or tenderness).
      - (b) The skeletal-muscle related symptoms occurred while receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or combination products); AND
      - (c) When receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or as combination products) the skeletal-related muscle symptoms resolved upon discontinuation of each respective statin therapy (atorvastatin and rosuvastatin); OR  
Note: Examples of skeletal-related muscle symptoms include myopathy and myalgia.
- B) Patient Currently Receiving Nexletol. Approve if according to the prescriber, the patient has experienced a response to therapy.  
Note: Examples of a response to therapy include decreasing low-density lipoprotein cholesterol (LDL-C), total cholesterol, non-high-density lipoprotein (non-HDL-C), or apolipoprotein B levels. Also, if the patient is currently receiving the requested therapy but has not previously received approval of Nexletol for this specific indication through the Coverage Review Department, review under criteria for Initial Therapy. If the patient is restarting therapy with Nexletol, Initial Therapy criteria must be met.
2. **Heterozygous Familial Hypercholesterolemia (HeFH).**\* Approve for 1 year if the patient meets ONE of the following (A or B):
- A) Initial Therapy. Approve if the patient meets all of the following (i, ii, and iii):
    - i. Patient is  $\geq$  18 years of age; AND
    - ii. Patient meets one of the following (a, b, or c):
      - a) Patient has an untreated low-density lipoprotein cholesterol (LDL-C) level  $\geq$  190 mg/dL (prior to treatment with antihyperlipidemic agents); OR

- b) Patient has genetic confirmation of HeFH by mutations in the low-density lipoprotein receptor, apolipoprotein B, proprotein convertase subtilisin kexin type 9, or low-density lipoprotein receptor adaptor protein 1 gene; OR
  - c) Patient has been diagnosed with heterozygous familial hypercholesterolemia meeting one of the following diagnostic criteria thresholds [(1) or (2)]:
    - (1) Prescriber confirms that the Dutch Lipid Network criteria score was > 5; OR
    - (2) Prescriber confirms that Simone Broome criteria met the threshold for “definite” or “possible (or probable)” familial hypercholesterolemia; AND
- iii. Patient meets one of the following (a or b):
- a) Patient meets all of the following [(1), (2), and (3)]:
    - (1) Patient has tried one high-intensity statin therapy (i.e., atorvastatin  $\geq$  40 mg daily; rosuvastatin  $\geq$  20 mg daily [as a single-entity or as a combination product]); AND
    - (2) Patient has tried one high-intensity statin above along with ezetimibe (as a single-entity or as a combination product) for  $\geq$  8 continuous weeks; AND
    - (3) LDL-C level after this treatment regimen remains  $\geq$  70 mg/dL; OR
  - b) Patient has been determined to be statin intolerant by meeting one of the following [(1) or (2)]:
    - (1) Patient experienced statin-related rhabdomyolysis; OR  
Note: Rhabdomyolysis is statin-induced muscle breakdown that is associated with markedly elevated creatine kinase levels (at least 10 times the upper limit of normal), along with evidence of end organ damage which can include signs of acute renal injury (noted by substantial increases in serum creatinine [Scr] levels [a  $\geq$  0.5 mg/dL increase in Scr or doubling of the Scr] and/or myoglobinuria [myoglobin present in urine]); OR
    - (2) Patient meets all of the following [(a), (b), and (c)]:
      - (a) Patient experienced skeletal-related muscle symptoms; AND  
Note: Examples of skeletal-related muscle symptoms include myopathy (muscle weakness) or myalgia (muscle aches, soreness, stiffness, or tenderness).
      - (b) The skeletal-muscle related symptoms occurred while receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or combination products); AND
      - (c) When receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or as combination products) the skeletal-related muscle symptoms resolved upon discontinuation of each respective statin therapy (atorvastatin and rosuvastatin); OR  
Note: Examples of skeletal-related muscle symptoms include myopathy and myalgia.
- B) Patient Currently Receiving Nexletol. Approve if according to the prescriber, the patient has experienced a response to therapy.  
Note: Examples of a response to therapy include decreasing low-density lipoprotein cholesterol (LDL-C), total cholesterol, non-high-density lipoprotein (non-HDL-C), or apolipoprotein B levels. Also, if the patient is currently receiving the requested therapy but has not previously received approval of Nexletol for this specific indication through the Coverage Review Department, review under criteria for Initial Therapy. If the patient is restarting therapy with Nexletol, Initial Therapy criteria must be met.

### Other Uses with Supportive Evidence

- 3. Primary Hyperlipidemia.\*** Approve for 1 year if the patient meets ONE of the following (A or B):  
Note: This is not associated with atherosclerotic cardiovascular disease (ASCVD) or heterozygous familial hypercholesterolemia (HeFH) and may be referred to as combined hyperlipidemia, hypercholesterolemia (pure, primary), dyslipidemia, or increased/elevated low-density lipoprotein cholesterol (LDL-C) levels.
- A) Initial Therapy.** Approve if the patient meets all of the following (i, ii, and iii):
- i.** Patient is  $\geq 18$  years of age; AND
  - ii.** Patient has a coronary artery calcium or calcification score  $\geq 300$  Agatston units; AND
  - iii.** Patient meets one of the following (a or b):
    - a)** Patient meets all of the following [(1), (2), and (3)]:
      - (1)** Patient has tried one high-intensity statin therapy (i.e., atorvastatin  $\geq 40$  mg daily; rosuvastatin  $\geq 20$  mg daily [as a single-entity or as a combination product]); AND
      - (2)** Patient has tried the one high-intensity statin therapy above along with ezetimibe (as a single-entity or as a combination product) for  $\geq 8$  continuous weeks; AND
      - (3)** LDL-C level after this treatment regimen remains  $\geq 100$  mg/dL; OR
    - b)** Patient has been determined to be statin intolerant by meeting one of the following [(1) or (2)]:
      - (1)** Patient experienced statin-related rhabdomyolysis; OR  
Note: Rhabdomyolysis is statin-induced muscle breakdown that is associated with markedly elevated creatine kinase levels (at least 10 times the upper limit of normal), along with evidence of end organ damage which can include signs of acute renal injury (noted by substantial increases in serum creatinine [Scr] levels [a  $\geq 0.5$  mg/dL increase in Scr or doubling of the Scr] and/or myoglobinuria [myoglobin present in urine]).
      - (2)** Patient meets all of the following [(a), (b), and (c)]:
        - (a)** Patient experienced skeletal-related muscle symptoms; AND  
Note: Examples of skeletal-related muscle symptoms include myopathy (muscle weakness) or myalgia (muscle aches, soreness, stiffness, or tenderness).
        - (b)** The skeletal-muscle related symptoms occurred while receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or combination products); AND
        - (c)** When receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or as combination products) the skeletal-related muscle symptoms resolved upon discontinuation of each respective statin therapy (atorvastatin and rosuvastatin); OR  
Note: Examples of skeletal-related muscle symptoms include myopathy and myalgia.
- B) Patient Currently Receiving Nexletol.** According to the prescriber, the patient has experienced a response to therapy.  
Note: Examples of a response to therapy include decreasing low-density lipoprotein cholesterol (LDL-C), total cholesterol, non-high-density lipoprotein (non-HDL-C), or apolipoprotein B levels. Also, if the patient is currently receiving the requested therapy but has not previously received approval of Nexletol for this specific indication through the Coverage Review Department, review under criteria for Initial Therapy. If the patient is restarting therapy with Nexletol, Initial Therapy criteria must be met.

**Note:**

\* A patient may have diagnoses that pertain to more than one indication, therefore, consider review under different approval conditions, if applicable (e.g., a patient with heterozygous familial

hypercholesterolemia may have had a clinical ASCVD event, a patient with primary hyperlipidemia may have heterozygous familial hypercholesterolemia).

### CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Nexletol is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

### REFERENCES

1. Nexletol® tablets [prescribing information]. Ann Arbor, MI: Esperion; December 2023.
2. Nissen SE, Lincoff AM, Brennan D, et al, for the CLEAR Outcomes investigators. Bempedoic acid and cardiovascular outcomes in statin-intolerant patients. *N Engl J Med*. 2023;388(15):1353-1364.
3. Lloyd-Jones DM, Morris PB, Ballantyne CM, et al. 2022 ACC Expert Consensus Decision Pathway on the Role of Non-Statins Therapies for LDL-Cholesterol Lowering in the Management of Atherosclerotic Cardiovascular Disease Risk. *J Am Coll Cardiol*. 2022;80(14):1366-1418.
4. Stone NJ, Robinson J, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice guidelines. *J Am Coll Cardiol*. 2014;63(25 Pt B):2889-2934.
5. Grundy SM, Stone NJ, Bailey AL, et al. 2018 ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol. A report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2019;139(35):e1082-1143.
6. Newman CB, Blaha MJ, Boord JB, et al. Lipid management in patients with endocrine disorders: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2020;105(12):3613-3682.
7. Handelsman Y, Jellinger PS, Guerin CK, et al. Consensus Statement by the American Association of Clinical Endocrinologists and the American College of Endocrinology on the management of dyslipidemia and prevention of cardiovascular disease algorithm-2020 executive summary. *Endocr Pract*. 2020;26(10):1196-1124.
8. Jacobson TA, Ito MK, Maki KC, et al. National Lipid Association recommendations for patient-centered management of dyslipidemia: Part 1-full report. *J Clin Lipidol*. 2015;9:129-169.
9. Gidding SS, Champagne MA, de Ferranti SD, et al. The agenda for familial hypercholesterolemia. A scientific statement from the American Heart Association. *Circulation*. 2015;132(22):2167-2192.
10. Banach M, Penson PE, Farnier M, et al. Bempedoic acid in the management of lipid disorders and cardiovascular risk. 2023 position paper of the International Lipid Expert Panel (ICEP). *Prog Cardiovasc Dis*. 2023 Mar 7. [Online ahead of print].
11. Hect HS, Cronin P, Blaha M, et al. 2016 SCCT/STR guidelines for coronary artery calcium scoring of noncontrast noncardiac chest CT scans: A report of the Society of Cardiovascular Computed Tomography and Society of Thoracic Radiology. *J Thorac Imaging*. 2017;32(5):W54-S66.
12. Greenland P, Blaha MJ, Budoff MJ, et al. Coronary calcium score and cardiovascular risk. *J Am Coll Cardiol*. 2018;72(4):434-447.
13. Razavi AC, Agatston AS, Shaw LJ, et al. Evolving role of calcium density in coronary artery calcium scoring and atherosclerotic cardiovascular disease risk. *J Am Coll Cardiol Img*. 2022;15:1648-1662.
14. Lehker A, Mukherjee D. Coronary calcium risk score and cardiovascular risk. *Curr Vasc Pharmacol*. 2021;19(3):280-284.
15. Haase A, Goldberg AC. Identification of people with heterozygous familial hypercholesterolemia. *Curr Opin Lipidol*. 2012;23:282-289.

## **APPENDIX A**

### **Simon Broome Register Diagnostic Criteria.<sup>15</sup>**

LDL-C – Low-density lipoprotein cholesterol; LDL – Low-density lipoprotein; APOB – Apolipoprotein B; PCSK9 – Proprotein convertase subtilisin kexin type 9.

**APPENDIX B.**

**Dutch Lipid Network Criteria.<sup>9</sup>**

LDL-C – Low-density lipoprotein cholesterol; CAD – Coronary artery disease; LDLR – Low-density lipoprotein receptor; APOB – Apolipoprotein B; PCSK9 – Proprotein convertase subtilisin kexin type 9.