

## PRIOR AUTHORIZATION POLICY

- POLICY:** Immune Globulin – Cytogam Prior Authorization Policy
- Cytogam® (human cytomegalovirus immune globulin intravenous infusion – Saol Therapeutics)

**REVIEW DATE:** 12/14/2022

---

### OVERVIEW

Cytogam, a human cytomegalovirus (CMV) immune globulin intravenous (IGIV), is indicated for the **prophylaxis of CMV disease** associated with transplantation of kidney, lung, liver, pancreas, and heart.<sup>1</sup>

### Other Uses With Supportive Evidence

Maternal transmission of CMV to the fetus may occur at any time during gestation, leading to congenital CMV.<sup>2</sup> A study of 304 pregnant women with a primary CMV infection were offered CMV IGIV. In the therapy group, 157 women were treated with CMV IGIV low dose (100 mg/kg/infusion given once every month) or high dose (200 mg/kg/infusion given once every 2 weeks for up to 3 doses if needed). The trial demonstrated that 56% of patients without CMV IGIV vs. 30% of patients receiving CMV IGIV developed congenital CMV infection.

### POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Cytogam. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days. Because of the specialized skills required for evaluation and diagnosis of patients treated with Cytogam as well as the monitoring required for adverse events and long-term efficacy, approval requires Cytogam to be prescribed by or in consultation with a physician who specializes in the condition being treated.

**Automation:** None.

### RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Cytogam is recommended in those who meet one of the following criteria:

#### FDA-Approved Indication

1. **Prophylaxis of Cytomegalovirus Associated with Solid Organ Transplant.** Approve for 4 months if the medication is prescribed by or in consultation with a physician affiliated with a transplant center, hematologist, or an infectious disease physician.

#### Other Uses with Supportive Evidence

2. **Cytomegalovirus Associated with Pregnancy.** Approve for 6 months if the medication is prescribed by or in consultation with an infectious disease physician or an obstetrician-gynecologist.

12/14/2022

© 2022. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

**CONDITIONS NOT RECOMMENDED FOR APPROVAL**

Coverage of Cytogam is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

**REFERENCES**

1. Cytogam intravenous infusion [prescribing information]. Roswell, GA: Saol Therapeutics; October 2020.
2. Swanson E, Schleiss M. Congenital cytomegalovirus infection: New prospects for prevention and therapy. *Pediatr Clin North Am.* 2013;60:335-349.